





CENTRAL LINE DRESSING CHANGE

STANDARD WORK

1 HAND HYGIENE 		
2 ASSESS	<ul style="list-style-type: none"> redness? swelling? pain? fever? chills? damaged catheter? loose stitches? If so, STOP. Notify Provider. 	<p>Time Talent and Materials</p> <p>1 Nurse, 12 Minutes</p> <ul style="list-style-type: none"> <input type="checkbox"/> Clean gloves <input type="checkbox"/> Sterile gloves <input type="checkbox"/> Sutureless securement device <input type="checkbox"/> NS flush <input type="checkbox"/> Central line dressing kit: <ul style="list-style-type: none"> <input type="checkbox"/> Dressing <input type="checkbox"/> Sterile gauze <input type="checkbox"/> Needleless injector caps <input type="checkbox"/> Masks for care team <input type="checkbox"/> Chlorhexidine
3 EXPLAIN	<ul style="list-style-type: none"> Steps of the procedure How you're preventing infection 	
4 GATHER ITEMS	→	
5 PREP STERILE FIELD	<ul style="list-style-type: none"> Put on mask Open occlusive dressing Drop injection caps & gauze on field Open chlorhexidine Open sutureless securement device 	
6 HAND HYGIENE: PUT ON CLEAN GLOVES 		
7 PRIVACY/POSITION	<ul style="list-style-type: none"> Position patient for access Move linens from site If site is internal jugular or subclavian: Ask patient to turn head away from the catheter 	
8 REMOVE OLD	<ul style="list-style-type: none"> Peel dressing toward insertion until you can remove securement device. Use 4 alcohol pads to loosen securement anchor pads. Gently pull the old device away from the skin. 	
9 HAND HYGIENE: PUT ON STERILE GLOVES 		
10 SCRUB SITE	<ul style="list-style-type: none"> Hold catheter with sterile gauze Scrub with 2% chlorhexidine for 30 seconds Air dry completely. Allow 60+ seconds 	<p>Warning: Placing an occlusive dressing on wet skin increases the chance of a reaction.</p>
11 APPLY NEW	<ul style="list-style-type: none"> Use skin prep to protect the skin Connect the catheter to the securement device before adhering it to the skin Apply new sterile occlusive dressing Document the date, time and your initials on the new dressing 	
12 FLUSH-CLAMP-REMOVE	<ul style="list-style-type: none"> Change needleless injection caps To keep blood out of the catheter when not in use, clamp the port before removing the syringe 	
13 ASSESS	<ul style="list-style-type: none"> Assess, treat and reassess pain if present 	
14 DISCARD	<ul style="list-style-type: none"> Discard used supplies 	
15 HAND HYGIENE 		
16 DOCUMENT	<ul style="list-style-type: none"> Document procedure in the electronic record 	